

ZAŠTITA OD BOLESTI




FUNGICHEL

Za preventivno suzbijanje **plamenjače** (*Phytophthora* spp.) i **paleži lista** (*Alternaria* spp., *Septoria* spp.) u količini od 1,5 l/ha u vrijeme kada su uvjeti za pojavu bolesti povoljni, svakih 15-20 dana

KORIJENASTO POVRĆE (MRKVA, PERŠIN, CELER)

PRIHRANA BIOSTIMULATORIMA

SADNJA	VEGETATIVNI PORAST	RAZVOJ KORIJENA	NALIJEVANJE KORIJENA
 <p>Odmah nakon presađivanja folijarno:</p> <p>STIMCAP 2,0 l/ha + CITOQUIN MIX 1,5 l/ha</p> <p>Nakon 10-15 dana folijarno:</p> <p>STIMCAP 2,0 l/ha + CITOQUIN MIX 1,5 l/ha + GROWCAP 3,0-4,0 l/ha</p>	 <p>folijarno</p> <p>CITOQUIN MIX 2,0 l/ha + GROWCAP 3,0-4,0 l/ha</p>	 <p>folijarno</p> <p>CITOQUIN MIX 2,0 l/ha + TRIXBOR 2,0 l/ha</p>	 <p>folijarno</p> <p>DELTAQUIN 2-3 puta svakih 15 dana 2,0 l/ha + GROWCAP 3,0 l/ha (PO POTREBI)</p>
			