

ZAŠTITA OD BOLESTI



FUNGICHEL

Za preventivno suzbijanje plamenjače (*Plasmopara viticola*) u koncentraciji od 0,1-0,15% (100-150 ml/100 lit vode), kretanjem vegetacije svakih 15-20 dana.

BEZ
KARENCE



BOTRYNE

Za preventivno i kurativno suzbijanje sive plijesni (*Botrytis cinerea*), u koncentraciji 0,25% (250 g/100 lit vode) nakon cvatnje 3-4 puta svakih 10-15 dana.

BEZ
KARENCE

VINOVA LOZA

PRIHRANA BIOSTIMULATORIMA

RAST MLADICA	CVATNJA	ZAMETANJE PLODOVA	RAZVOJ PLODOVA 8-10 mm	DOZRIJEVANJE
 <p>MLADICE 8-10 cm CITOQUIM MIX 0,25% (250 ml/100 lit vode) + GROWCAP 0,3% (300 ml/100 lit vode)</p>	 <p>PRED CVATNJU CITOQUIM MIX 0,25% (250 ml/100 lit vode) + TRIXBOR 0,2% (200 ml/100 lit vode)</p>	 <p>TRIXBOR 0,2% (200 ml/100 lit vode)</p>	 <p>BOBICE 8-10 mm DELTAQUIN 0,25% (250 ml/100 lit vode) + ENERMAX 0,075% (75 g/100 lit vode) + ACTIV Ca 0,3% (300 ml/100 lit vode)</p>	 <p>POČETKOM ŠARANJA BOBICA CITOQUIM MIX 0,25% (250 ml/100 lit vode) + GROWCAP 0,3% (300 ml/100 lit vode)</p>

